

# Human-sloth bear conflict: Causes and Mitigation in Kota, Rajasthan, India

*The sloth bear (*Melursus ursinus*), is the most dangerous wild animal due to its' unpredictable behaviour and often attacks on human in forests. The present investigation represents human-sloth bear conflicts, as per standard methods and survey. The results suggested that, most of the casualties can be reduced by restricting human entries into sloth bear habitats, avoiding camping and housing in their habitats especially near water sources. Further road accidents were common in the study area due to National Highway near the forest area. The scientific management in collaboration with local inhabitants will support the conservation of this species.*

**Key words:** *Melursus ursinus*, Conflicts, Endangered, Interviews, Forests.

## Introduction

Wildlife conservation has always been challenging task in India as there is a need to uphold a balanced approach between natural resource conservation and meeting the growing needs of human population and development. The basic requirement to evolve strategies for conservation of endangered species and their natural habitats on one hand and on the other hand they have to successfully deal with wildlife and Human conflicts. Although previous studies and literature revealed that Human and wildlife have a long time co-existence. Due to the competition for resources at different levels human-wildlife conflicts increased manifold. They typically compete directly with people for resources, space, food, security and cover. All bear species can kill or injure livestock, damage agricultural or horticulture crops, or otherwise directly compete with people (Charoo *et al.*, 2009).

The sloth bear is endemic to the Indian subcontinent, having geographical distribution across India, Nepal and Sri Lanka. The species has been extirpated from Bangladesh and is reported to be rare in Bhutan (Garshelis *et al.*, 2008). The sloth bear's range in India extends from the foothills of the Himalayas to the southern tip of the Western Ghats; however, its distribution is non-continuous and fragmented. The central Indian highlands, Western Ghats and the Eastern Ghats are considered to be strongholds of the sloth bear (Jhala *et al.*, 2011 and Yoganand *et al.*, 2013); central India harbors the largest intact habitat and population of bears (Sathyakumar *et al.*, 2012). In India, sloth bear occur frequently in moist and dry deciduous forest (42% and 33% respectively) and less frequently in wet evergreen (13%) and dry scrub (6%) forests (Yoganand *et al.*, 2006). In India sloth bears are reported to exist in 14 protected areas including 46 National Parks and 128 Wildlife Sanctuaries (Chauhan, 2006).

The Chambal River passes through a deep gorge created by a great boundary fault in the Vindhya on the Hadauti Plateau of Rajasthan. The upper tier plateau land from the river bed ranges from 60.5 m. The river flowing through the gorge drops 87m. Along a length of 48km three dams and barrage have changed the flow of this river, which was once a torrentially free flowing stream as it poured through the deep gorge. The Chambal River valley was strong hold of Big Cats in the pre-independence era, possibly the only area where trigger and leopard hunts were routinely

*Human-sloth bear conflicts is represented as per standard methods and survey, with five years records which were documented from March, 2010 to March, 2016.*

**KIRAN CHOUDHARY AND  
KRISHNENDRA SINGH NAMA<sup>1</sup>**  
M.B. (PG) College, Kota (Rajasthan)  
E-mail: choudharykiran01@gmail.com;  
namasahib@gmail.com

Received May, 2017  
Accepted February, 2018

<sup>1</sup>Career Point University Kota (Rajasthan)

organised from boats. The flat plateau land over the gorge is interspersed with dry deciduous forest of *Anogeissus pendula* (Dhonk) and *Boswellia serrata* (Salar). The plateau land near Kota is stony scrub land with scattered *Zizyphus nummularia* (Ber) and *Capparis decidua* and *Capparis sepiaria* (Jal) bushes, which explains why bears are so plentiful. The vistas, generally sparse and the vegetation is coarse. The glimpse into the valley offers a sticking contrast. The valley is evergreen with a profusion of trees, bushes and semi aquatic grasses. The stretches of bamboo also cloth the valley floor with intermediate outcroppings of *Sterculia urens* (Gumkaraya), *Mitragyna parviflora* (Kalam), *Terminalia arjuna* (Arjun), *Lannea coromandelica* (Gurjan), *Cassia fistula* (Amaltash), etc.

The topography of the valley continues to be ideal for several species that have learned to use its ledges, nooks and rocky crannies to their advantages. Long billed vultures, Egyptian vultures, brown fish owls, dusky eagle owl and Indian eagle owl all breed here. The bears of the Chambal have Leopards, Hyena, Porcupine, Blue Bull, Chinkara, langur and an occasional wolf for company. Sometimes any illusive Fishing Cat or a family of Smooth Indian Otters may be encountered at the water edges. Present study was aimed to study the incidence of road accident, human attacks and ingress of bears in areas of human settlement/ outside the forest areas and the mitigation measurements of Human-Sloth Bear conflicts.

## Methodology

To study the Bear-Human conflict in the study area following methodology were chosen-

### Field survey

Regular field visits were done to determine sloth bear presence or absence and to study its ecology and behaviour in the present study area from 2010-2016.

### Questionnaire survey

Informal interviews using semi-structured questionnaires were carried out in affected the villages located on and beyond the periphery of forest areas of Kota district.

In the first phase of survey five villagers were chosen randomly from each exaggerated village. They were interviewed to record the frequency of conflicts, probable causes and local desire for easing of attacks.

In the second phase of survey those people were interviewed personally who faced sloth bear attack, so they were the real victims and eye witness. It'll helpful to assess the circumstances during attack, and in formulations of probable management tactics.

### Record survey

Data allied to aspects such as road accidents of sloth bear were intensively surveyed. Six year records were documented since March 2010 to March 2016 so the accidental prone zone can be identified around the study area.

Review of previous studies which were conducted in earlier decade were also studied to comprehend the behavioural ecology of sloth bear in various situations viz. in wild and while encountered with human.

## Personal involvement

Involvement in rescue operation of sloth bears with forest department were the most effective method to predict the behaviour of bear and contemplations of local people for these sort of wildlife encounters.

## Observations

### Ecology and Behaviour

Sloth bear, forested animal found in areas with dense canopy cover where human disturbance is limited or restricted. It is typically semi-nocturnal, although occasionally it is active during day time. Post monsoon when food is aplenty across a larger landscape, sloth bears tend to leave the Chambal river valley, only to return in the summer when their favourite trees *Zizyphus sp.*, *Madhuca indica* (Mahua), *Syzygium cumini* (Jamun) etc. are in fruit, bee hive are available to raid and termite mounds are stocked with insects, food that relish bears. During an observation any encounters between the sloth bears and leopards, hyenas or wolves were not observed (Vyas *et al.*, 2014). This was an account of the sheer abundance of food and relatively low population densities of carnivores.

Interestingly Bears and Langoors living in close proximity displayed no hostility towards each other at all. Nor did pea fowl seem to bother the bears, thought they did always maintain a safe flight distance.

The reason for the thriving sloth bear population the Chambal area is hardly surprising. The area is rich in forage, with an abundance of termite mounds and *Zizyphus jujuba* bushes on the plateau. Honey combs are plentiful in the valley, which also offers a reliable source of water, but most critically, the area enjoys protected status and humans and animals are not allowed to compete for resources. There is little more that sloth bears, all most other wild animals for that matter, really need to survive.

Over the past 5 yrs., sightings of bears have increased palpably, with multiple groups of adults, and sub adults occupying the habitats. In winter, solitary adults and sub adults can be seen all through the day, scouring the plateau for food. In summer, family seemed to be closely knit and one can able to spot them only at dawn and dusk.

### Sloth Bear-Human Conflicts

Intensive five year's study and Forest (Wildlife) Department documents illustrates that Sloth bear conflicts are occasional in this area (Fig.1). According to the preliminary studies the possible causes for these incidences of Human-Sloth bear conflicts are-

1. Hustle vehicular movement on National Highway-76 in close proximity of bear habitat is a ground cause for road accidents.
2. Shrinking habitat due to extension of agricultural lands and anthropogenic encroachment are the core causes of habitat degradation.
3. Increased dependency on natural resources leading to increased frequency of bear-human encounters.
4. Increasing human and livestock population in and around the Protected Areas and forest areas through a direct pressure on sloth bear movement.



Fig. 1: Map showing study area.



Fig. 2: Road accident.

### Conflict types

#### Road accidents

National Highway-76 via Kota city can be considered as a channel between Rajasthan to Gujrat states. So it has a huge traffic burden especially in peak hours of night. In a very close proximity of highway Jawahar Sagar Sanctuary and National Chambal Sanctuaries are situated, which are cradle home for Sloth Bears since long. As a habit bears wander for their food mainly at night and cross this highway regularly. A speedy vehicle passes through the highway and strike with these animals is the main cause of unintended death (Fig. 2).

#### Human attacks

Increasing human population is directly proportional to increased dependence on forests resources; as well as it leads increased frequency of conflicts of two mammalian species viz. Human and sloth bear in and around the forest area. In the present study area villagers move inside the forest boundary legally and illegally to collect fuel wood and for mining purpose. Hence they are considered as disturbing elements by sloth bear in their close vicinity. Bear generally doesn't attack people unless and until, it feels threatened. Especially a female bear with her cubs offers a sudden encounter in the course of their protection. This might be fatal for any one in front of her.



## ***Ingression of Sloth bear inside the City***

Sloth bear are voracious feeder as they have great sense of smell they are attracted by cuisines prepared nearby human settlements of sloth bear habitat. It offers sloth bear doorway inside the human residential areas (Fig. 3)

## **Result and Discussion**


Field work and questionnaire survey was conducted to prepare final findings regarding human-sloth bear conflicts in study area. Almost all the villages within that area is surveyed and villagers from each village were interviewed to know their perception towards the presence of bear in the area. Survey was conducted in 19 villages of study area to inquire about sloth bear. Approximately 71% of interviewees reported that they had

regular sightings of sloth bears on their farms and nearby water sources. About 25% of the interviewees reported that they have not seen sloth bears in nearby areas and 4% were uncertain about the presence or absence of bears in the adjoining areas.

The bear ingression in nearby villages were also analysed on the basis of time of attack, which shows that majority of incidents occurred during the day time. Further it was also recorded that maximum conflicts occurred in winter season (Table 1). Winter is the fruiting season of most of the plants occurring in the study area, these fruits are consumed by sloth bears. Result shows that maximum conflicts are in the form of road accidents (Fig. 4).



**Fig. 3:** Sloth bear rescue.



**Fig. 4:** Per cent occurrence of different conflicts in the study area.

**Table 1:** Occupancy of conflicts in different months in different areas.

Site of Incidence	Month of incidence	Ingression rate	Road accidents	Human attacks
Khadipur	Aug. 2009	-	2	--
Khadipur	Oct. 2009	-	1	-
Karondi	June 2011	-	2	-
KTPS	July 2013	1	-	-
Karmi Mata Temple	Oct. 2013	1	-	-
Nanta	Dec. 2013	1	-	-
Bhainsroadgarh	Feb. 2014	-	1	-
KTPS	April 2014	1	-	-
Nanta	Dec. 2014	1	-	-
Doliya	Mar. 2016	-	-	1
Ganeshpura	May 2016	-	1	-

### Probable management practices for human-sloth bear conflicts

Based on questionnaire and field surveys carried out in the 7 villages of 3 ranges of Jawahar sagar sanctuary, some basic understanding of the human-sloth bear conflicts in this region has been obtained. Some of the preliminary recommendations for understanding and mitigating human-sloth bear conflicts are given below.

#### Creation of conflict management team

A fully equipped, well trained and motivated 'Conflict Management Team' comprising of wildlife staff, veterinarians, staff of related line departments or institutions, and wild life NGOs has to be formed at the wild life division level to respond to conflict situations, including bear rescue, treatment, translocation and monitoring.

#### Control rooms

At district level control rooms and management teams should be established by the department for handling wildlife-human conflict cases.

#### Awareness creation

Knowledge of sloth bear behaviour is essential for the villagers who live near bear habitats and are vulnerable to bear attacks. Awareness creation through elected bodies at village or block levels, religious or educational establishments, and other government or non-governmental agencies needs to be explored. Simple precautionary measures such as avoiding moving singly during dawns and dusks in crop fields or forests during summer and autumn could reduce bear attacks on humans.

#### Monitoring of Conflict areas

Use of passive and preventive actions is an important step in conflict management. There should be proper monitoring of sloth bear-human conflict areas where at least one of the employees is present at the problem site until the situation is resolved. While monitoring the bear, information such as number, age, location, bear behaviour and the direction of the movement of the conflicting animal has to be recorded and reported. Movement of the people should be restricted to give safe passage to the bear.

#### Translocation and marking of problem bear

The problem bear that are captured and translocated from the conflict areas should be marked by radio collar, fluorescent collar or by ear tags prior to their release back into the wild so that the movement and the ranging patterns of these animals could be studied. This will be very helpful in better understanding of problem bears and ultimately help in mitigating the problem.

#### Fencing and Underpass

NH-76 is the major problem for the life of wild animals as it is a lethal highway. Most of the accidents occur in the morning hours and in late evening as many animals are nocturnal and they left their home for the feeding purpose and got accidental death. So there should be fencing on both the sides of Highway and if possible there should be a protocol of underpass in and around the forest areas.

#### Other practices

Signage boards on roads regarding speed control should be present on the highways. Slogans on forest walls indicating importance of wildlife can aware peoples regarding the safety of these wild animals.

#### Conclusion

Present study reveals that Sloth Bear-Human conflict in the form of human attack, crop loss and livestock damage is very low but on the other hand death of wild animals in road accidents is very high mostly in rainy and winter season. But well-designed human-wildlife conflict management plans which amalgamate different techniques and are adjusted based on the nature of predicament can heighten co-existence. Impending solutions should be well thought-out and selected based on their efficacy, cost and adequacy by the society. These practices can reduce the negative attitudes of communities towards wildlife and preservation.

The abundance of this species has decreased since the 1950s to <10,000 bears in 1998. This decrease was due to increased human population, which have altered forests and harassed bears (Cowan, 1972). The sloth bear is now classified as endangered by the Government of India (Wildlife Protection Act 1972 Amended). The sloth bear is an omnivorous animal. Habitats of sloth bear has destructed and degrading continuously as a result sloth bears have been extirpated in some areas of India (Singh, 2001). Meanwhile humans compete directly with bears by consuming the bears' food resources (Rajpurohit, 2000).

कोटा, राजस्थान, भारत में मानव-रीछ संघर्ष : कारण एवं  
न्यूनीकरण

किरण चौधरी और कृष्णोन्द्र सिंह नामा

सारांश

रीछ ( *मीलूसर्स यूर्सिनस* ) अपने अप्रत्याशित व्यवहार के कारण सबसे खतरनाक जंगली पशु है तथा प्रायः जंगलों में मानवों पर आक्रमण कर देता है। वर्तमान अन्वेषण में मानक विधियों एवं सर्वेक्षण के अनुसार मानव-रीछ संघर्ष का प्रतिनिधित्व है। परिणाम संकेत करते हैं कि अधिकांश दुर्घटनाओं के रीछ आवासों में मानव प्रवेश को प्रतिबंधित करके तथा इनके आवासों में विशेषकर पानी के स्रोत के समीप कैम्पिंग तथा हाउसिंग का बचाव करके कम किया जा सकता है। इसके अलावा वन क्षेत्रों के नजदीक राष्ट्रीय राजमार्ग के कारण अध्ययन क्षेत्र में सड़क दुर्घटनाएं आम थी। स्थानीय निवासियों के सहयोग से वैज्ञानिक प्रबंधन इस प्रजाति के संरक्षण में सहायता करेंगे।

## References

Charoo S.A., Sharma L.K. and Sathyakumar S. (2009). Asiatic Black Bear-Human Conflicts around Dachigam National Park, Kashmir. Technical Report. Wildlife Institute of India, Dehradun 51pp.

Chauhan N.P.S. (2006). The Status of sloth bear in India. In Understanding Asian bears to secure their future, Japan Bear Network, Ibaraki, Japan. 26-34 pp.

Cowan I.Mc.T. (1972). The Status and Conservation of bears (Ursidae) of the world-1970. *Inter. Conference on Bear Research and Management*. 2 342-367.

Garshelis D.L., Ratnayake S. and Chauhan N.P.S. (2008). (IUCN SSC Bear Specialist Group). c2008 [cited 2 July 2016]. In: *Melursus ursinus*. The IUCN Red List of Threatened Species: e T13143A3413440.

[http://projecttiger.nic.in/WriteReadData/PublicationFile/Tiger\\_Status\\_oct\\_2010.pdf](http://projecttiger.nic.in/WriteReadData/PublicationFile/Tiger_Status_oct_2010.pdf)

<http://www.iucnredlist.org/details/summary/13143/0>

Jhala Y.V., Qureshi Q., Gopal R. and Sinha P.R. (2011). Status of tigers, co-predators and prey in India. New Delhi and Dehradun: National Tiger Conservation Authority, Government of India, New Delhi, and Wildlife Institute of India. Report No. TR 2011/003.

Rajpurohit K.S. and Krausman P.M. (2000). Huan-sloth bear conflicts in Madhya radish, India. *Wildlife Society Bulletin*, 28 393-399.

Sathyakumar S., Kaul R., Ashraf N.V.K., Mookerjee A. and Menon V. (2012). (Ministry of Environment and Forests, Government of India Wildlife Institute of India, Wildlife Trust of India, Central Zoo Authority, Government of India). National bear conservation and welfare action plan; 2012. Technical report. India: Ministry of Environment and Forests, Wildlife Institute of India and Wildlife Trust of India.

Singh H.S. (2001). Natural heritage of Gujarat. Gujarat Ecological Education and Research Foundation, Gandhinagar, India. 262 pp.

Vyas R., Tomar R.S. and Arya M. (2014). Call of the Wild Sanctuary Asia Bear necessities. *Hornbill Article*. 84-87 pp.

Yoganand K., Rice C.G., Johnsing A.J.T. and Seidensticker J. (2006). Is the sloth bear in India secure? A preliminary report on distribution, threats and conservation requirements. *J. Bombay Nat. Hist. Soc.*, 103(2-3): 172-181.

Yoganand K., Rice C.G. and Johnsingh A.J.T. (2013). Sloth bear: *Melursus ursinus*. In: *Mammals of South Asia* (Johnsingh AJT, Manjrekar N, editors) Universities Press. 1: 438-456.

## Acknowledgements

Authors are thankful to the team members for doing research and questionnaire survey in the villages. They are also greatly thankful to Forest and wildlife department of Kota for official support, facilities and encouragement during the research. Contribution of Mr. Ravindra Singh Tomar and Mr. Manish Arya is also unforgettable for providing important information.